

Belimumab (Benlysta)

Provider Order Form rev. 1/31/23



14107 Cortez Blvd
Brooksville FL 34613
P: (352)549-9962
F: (352)549-9963
www.360infusioncenter.com
E: info@360infusioncenter.com

PATIENT INFORMATION

Referral Status: New Referral Updated Order Order Renewal

Date: _____ Patient Name: _____ DOB: _____

ICD-10 code (required): _____ ICD-10 description: _____

NKDA Allergies: _____ Weight (lbs/kg): _____ Height: _____

Patient Status: New to Therapy Continuing Therapy Last Treatment Date: _____ Next Due Date: _____

PROVIDER INFORMATION

Referral Coordinator Name: _____ Referral Coordinator Email: _____

Ordering Provider: _____ Provider NPI: _____

Referring Practice Name: _____ Phone: _____ Fax: _____

Practice Address: _____ City: _____ State: _____ Zip Code: _____

NURSING

Provide nursing care per 360 Infusion Center Nursing Procedures, including reaction management and post-procedure observation

LABORATORY ORDERS

- CBC at each dose every _____
- CMP at each dose every _____
- CRP at each dose every _____
- Other: _____

PRE-MEDICATION ORDERS

- acetaminophen (Tylenol) 500mg / 650mg / 1000mg PO
- cetirizine (Zyrtec) 10mg PO
- loratadine (Claritin) 10mg PO
- diphenhydramine (Benadryl) 25mg / 50mg PO / IV
- methylprednisolone (Solu-Medrol) 40mg / 125mg IV
- hydrocortisone (Solu-Cortef) 100mg IV
- Other: _____
- Dose: _____ Route: _____
- Frequency: _____

THERAPY ADMINISTRATION

- Belimumab (Benlysta) in 250ml 0.9% sodium chloride, intravenous infusion over one hour
 - Dose: 10mg/kg
 - Route: 0 intravenous
 - Frequency: induction: week 0, 2, 4, and then every 4 weeks / maintenance: every 4 weeks / other: _____
 - Infuse over one hour
 - Flush with 0.9% sodium chloride at infusion completion

- Belimumab (Benlysta) injection
 - Dose: 200mg
 - Frequency: once weekly
 - Route: 0 subcutaneous

- Patient is required to stay for 30-minute observation
- Refills: Zero / for 12 months / _____
(if not indicated order will expire one year from date signed)

SPECIAL INSTRUCTIONS

*Consider administering premedication for prophylaxis against infusion reactions and hypersensitivity reactions.

Provider Name (Print)

Provider Signature

Date