

# Certolizumab (Cimzia)

Provider Order Form rev. 10/12/2022



14107 Cortez Blvd  
Brooksville FL 34613  
P: (352)549-9962  
F: (352)549-9963  
[www.360infusioncenter.com](http://www.360infusioncenter.com)  
E: info@360infusioncenter.com

## PATIENT INFORMATION

Referral Status:  New Referral  Updated Order  Order Renewal

Date: \_\_\_\_\_ Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_

ICD-10 code (required): \_\_\_\_\_ ICD-10 description: \_\_\_\_\_

NKDA Allergies: \_\_\_\_\_ Weight (lbs/kg): \_\_\_\_\_ Height: \_\_\_\_\_

Patient Status:  New to Therapy  Continuing Therapy Last Treatment Date: \_\_\_\_\_ Next Due Date: \_\_\_\_\_

## PROVIDER INFORMATION

Referral Coordinator Name: \_\_\_\_\_ Referral Coordinator Email: \_\_\_\_\_

Ordering Provider: \_\_\_\_\_ Provider NPI: \_\_\_\_\_

Referring Practice Name: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_

Practice Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip Code: \_\_\_\_\_

## NURSING

- Provide nursing care per 360 Infusion Center Nursing Procedures, including reaction management and post-procedure observation
- TB status & date (list results here & attach clinicals)  
\_\_\_\_\_
- Hepatitis B status & date (list results here & attach clinicals)  
\_\_\_\_\_

## THERAPY ADMINISTRATION

- Certolizumab** (Cimzia) subcutaneous injection
  - Induction
    - Dose:  400mg /  \_\_\_\_\_mg at Week 0, 2, 4, and then with maintenance dosing below
    - Give each 200mg injection separately
  - Maintenance
    - Dose:  200mg /  400mg
    - Frequency:  every 2 weeks /  every 4 weeks /  other: \_\_\_\_\_
- Patient is required to stay for 30-minute observation
- Refills:  Zero /  for 12 months /  \_\_\_\_\_  
(if not indicated order will expire one year from date signed)

## PRE-MEDICATION ORDERS

- acetaminophen (Tylenol)  500mg /  650mg /  1000mg PO
- cetirizine (Zyrtec) 10mg PO
- loratadine (Claritin) 10mg PO
- diphenhydramine (Benadryl)  25mg /  50mg  PO /  IV
- methylprednisolone (Solu-Medrol)  40mg /  125mg IV
- hydrocortisone (Solu-Cortef)  100mg IV
- Other: \_\_\_\_\_  
Dose: \_\_\_\_\_ Route: \_\_\_\_\_  
Frequency: \_\_\_\_\_

## SPECIAL INSTRUCTIONS

\*Evaluate patients for tuberculosis risk factors and test for latent infection prior to initiating CIMZIA and periodically during therapy. Treatment of latent tuberculosis infection prior to therapy with TNF-blocking agents has shown to reduce risk of tuberculosis reactivation during therapy. Prior to initiating CIMZIA, assess if treatment for latent tuberculosis is needed; and consider an induration of 5 mm or greater a positive tuberculin skin test result, even for patients previously vaccinated with Bacille Calmette-Guerin (BCG).

\*Test patients for HBV infection before initiating treatment with CIMZIA. For patients who test positive for HBV infection, consultation with a physician with expertise in the treatment of hepatitis B is recommended.

Provider Name (Print) \_\_\_\_\_ Provider Signature \_\_\_\_\_ Date \_\_\_\_\_

