

# Alglucosidase alfa (Lumizyme)

Provider Order Form rev. 3/1/2023



14107 Cortez Blvd  
Brooksville FL 34613  
P: (352)549-9962  
F: (352)549-9963  
[www.360infusioncenter.com](http://www.360infusioncenter.com)  
E: info@360infusioncenter.com

## PATIENT INFORMATION

Referral Status:  New Referral  Updated Order  Order Renewal

Date: \_\_\_\_\_ Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_

ICD-10 code (required): \_\_\_\_\_ ICD-10 description: \_\_\_\_\_

NKDA Allergies: \_\_\_\_\_ Weight (lbs/kg): \_\_\_\_\_ Height: \_\_\_\_\_

Patient Status:  New to Therapy  Continuing Therapy Last Treatment Date: \_\_\_\_\_ Next Due Date: \_\_\_\_\_

## PROVIDER INFORMATION

Referral Coordinator Name: \_\_\_\_\_ Referral Coordinator Email: \_\_\_\_\_

Ordering Provider: \_\_\_\_\_ Provider NPI: \_\_\_\_\_

Referring Practice Name: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_

Practice Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip Code: \_\_\_\_\_

## NURSING

- Provide nursing care per 360 Infusion Center Nursing Procedures, including reaction management and post-procedure observation

## LABORATORY ORDERS

- CBC  at each dose  every \_\_\_\_\_
- CMP  at each dose  every \_\_\_\_\_
- CRP  at each dose  every \_\_\_\_\_
- Other: \_\_\_\_\_

## PRE-MEDICATION ORDERS

- acetaminophen (Tylenol)  500mg /  650mg /  1000mg PO
- cetirizine (Zyrtec) 10mg PO
- loratadine (Claritin) 10mg PO
- diphenhydramine (Benadryl)  25mg /  50mg  PO /  IV
- methylprednisolone (Solu-Medrol)  40mg /  125mg IV
- hydrocortisone (Solu-Cortef)  100mg IV
- Other: \_\_\_\_\_
- Dose: \_\_\_\_\_ Route: \_\_\_\_\_
- Frequency: \_\_\_\_\_

## THERAPY ADMINISTRATION

- Alglucosidase alfa** (Lumizyme) in 0.9% sodium chloride, intravenous infusion, final concentration of 0.5 to 4mg/ml, administer with 0.2 micron filter
  - Dose:  20mg/kg /  other \_\_\_\_\_
  - Frequency:  every 2 weeks
  - Administer over approximately 4 hours, in a step wise manner. Initial infusion rate should be no more than 1mg/kg/hr. Infusion rate may be increased by 2mg/kg/hr every 30 minutes after patient tolerance is established. Max rate is 7mg/kg/hr. If the patient is stable, alglucosidase alfa may be administered at the maximum rate of 7mg/kg/hr until the infusion is completed
- Flush with 0.9% sodium chloride at infusion completion
- Patient is required to stay for 30-minute observation period
- Refills:  Zero /  for 12 months /  \_\_\_\_\_  
(if not indicated order will expire one year from date signed)

## SPECIAL INSTRUCTIONS

Provider Name (Print)

Provider Signature

Date