

Teprotumumab-trbw (Tepezza)

Provider Order Form rev. 3/1/2023



14107 Cortez Blvd
Brooksville FL 34613
P: (352)549-9962
F: (352)549-9963
www.360infusioncenter.com
E: info@360infusioncenter.com

PATIENT INFORMATION

Referral Status: New Referral Updated Order Order Renewal

Date: _____ Patient Name: _____ DOB: _____

ICD-10 code (required): _____ ICD-10 description: _____

NKDA Allergies: _____ Weight (lbs/kg): _____ Height: _____

Patient Status: New to Therapy Continuing Therapy Last Treatment Date: _____ Next Due Date: _____

PROVIDER INFORMATION

Referral Coordinator Name: _____ Referral Coordinator Email: _____

Ordering Provider: _____ Provider NPI: _____

Referring Practice Name: _____ Phone: _____ Fax: _____

Practice Address: _____ City: _____ State: _____ Zip Code: _____

NURSING

- Provide nursing care per 360 Infusion Center Nursing Procedures, including reaction management and post-procedure observation

LABORATORY ORDERS

- CBC at each dose every _____
- CMP at each dose every _____
(CMP includes serum blood glucose)
- Other: _____

PRE-MEDICATION ORDERS

- acetaminophen (Tylenol) 500mg / 650mg / 1000mg PO
- cetirizine (Zyrtec) 10mg PO
- loratadine (Claritin) 10mg PO
- diphenhydramine (Benadryl) 25mg / 50mg PO / IV
- methylprednisolone (Solu-Medrol) 40mg / 125mg IV
- hydrocortisone (Solu-Cortef) 100mg IV
- Other: _____
Dose: _____ Route: _____
Frequency: _____

THERAPY ADMINISTRATION

- Teprotumumab-trbw** (Tepezza) in 0.9% sodium chloride, intravenous infusion
 - Dose: (Indicate if patient has received any previous doses.)
 - 10mg/kg for the first infusion
 - 20mg/kg for infusions 2-8
 - Frequency: Every 3 weeks, 8 total infusions.
 - Administer the first 2 infusions over 90min. Subsequent infusions may be reduced to 60min if well tolerated. If reaction occurs, interrupt or slow the rate of infusion.
 - Dilute with 0.9% Sodium Chloride. For doses <1800mg use a 100ml bag. For doses ≥1800mg use a 250ml bag. (Remove equal volume.)
- Flush with 0.9% sodium chloride at the completion of infusion
- Patient is required to stay for 30-minute observation period
- Order is valid for 8 total infusions unless otherwise indicated.
(Order will expire one year from date signed)

SPECIAL INSTRUCTIONS

No premedication required. If the patient experiences an infusion reaction consider premedicating with an antihistamine, antipyretic, and/or corticosteroid.

TEPEZZA may cause an exacerbation of preexisting inflammatory bowel disease (IBD). Monitor patients with IBD for flare of disease.

Hyperglycemia or increased blood glucose may occur in patients treated with TEPEZZA. Monitor patients for elevated blood glucose and symptoms of hyperglycemia while on treatment with TEPEZZA. Patients with pre-existing diabetes should be under appropriate glycemic control before receiving TEPEZZA

Provider Name (Print)

Provider Signature

Date